






Atoms, Molecules and Ions




...Η πορεία του ατομικού προτύπου



Από την εποχή του Dalton μέχρι τον Schrödinger, το ατομικό μας πρότυπο τροποποιήθηκε πολλές φορές.


Η ΑΤΟΜΙΚΗ ΥΠΟΘΕΣΗ ΤΟΥ ΔΗΜΟΚΡΙΤΟΥ (400 π.Χ.)

- Η ύλη αποτελείται από μικροσκοπικά, αόρατα σωματίδια, τα οποία δεν μπορούν να διασπασθούν σε μικρότερα σωματίδια.
- Τα σωματίδια διαφέρουν ως προς το μέγεθος, το σχήμα και το βάρος.
- Υπάρχει κενός χώρος μεταξύ των σωματιδίων.
- Τα σωματίδια βρίσκονται σε συνεχή κίνηση μέσα στον κενό χώρο και συγκρούονται μεταξύ τους.
- Μετά τη σύγκρουση τα άτομα μπορούν να αποχωριστούν ή να ενωθούν.




ΙΣΤΟΡΙΚΗ ΑΝΑΔΡΟΜΗ

ΑΡΙΣΤΟΤΕΛΗΣ...
Απέρριψε την ατομική υπόθεση



... 2000 χρόνια (ΜΕΣΑΙΩΝΑΣ)...



Ατομική Θεωρία – John Dalton ~ 18


John Dalton Atomic Model

- Elements are made of extremely small particles called atoms
- Atoms of a given element are identical in size, mass, and other properties.
- Atoms cannot be subdivided, created, or destroyed
- Atoms of different elements combine in simple whole number ratios to form chemical compounds
- In chemical reactions, atoms are combined, separated, or rearranged.




Dalton's Atomic Theory


Carbon monoxide




Carbon dioxide




Law of Multiple Proportions



Atoms of element X



Atoms of element Y




Compounds of elements X and Y



Law of Conservation of Mass

ΙΣΤΟΡΙΚΗ ΑΝΑΔΡΟΜΗ

- **1897:** ανακάλυψη του ηλεκτρονίου (Thomson)



ΤΟ ΜΟΝΤΕΛΟ ΤΟΥ
«ΣΤΑΦΙΔΟΨΩΜΟΥ»




- **1911:** αναγνωρίζεται η ύπαρξη πυρήνα στο άτομο, ο οποίος αποτελεί το μεγαλύτερο ποσοστό της μάζας του, και είναι θετικά φορτισμένος (Rutherford)



- **1932:** ανακάλυψη του νετρονίου (Chadwick)



Cathode Ray Tube




J.J. Thomson, **measured mass/charge of e^-**

(1906 Nobel Prize in Physics) 9


Thomson's Model

Positive charge spread
over the entire sphere



Rutherford's Experiment

(1908 Nobel Prize in Chemistry)




1. atoms positive charge is concentrated in the nucleus
2. proton (p) has opposite (+) charge of electron (-)
3. mass of p is 1840 x mass of e^- (1.67×10^{-24} g)

ΑΤΟΜΙΚΟ ΠΡΟΤΥΠΟ Rutherford

- Τα άτομα έχουν πυρήνα, ο οποίος είναι θετικά φορτισμένος, καταλαμβάνει πολύ μικρό όγκο, αλλά περιέχει το μεγαλύτερο ποσοστό της μάζας του ατόμου (~99,5%).

- Τα ηλεκτρόνια περιφέρονται γύρω από τον θετικό πυρήνα.



ύρω από

ΔΟΜΗ ΤΟΥ ΑΤΟΜΟΥ

Πρωτόνιο

θετικά φορτισμένο σωματίδιο

$$m = 1,67252 \times 10^{-24} \text{ g}$$

$$q = 1,6022 \times 10^{-19} \text{ C}$$

Νετρόνιο

ηλεκτρικά ουδέτερο σωματίδιο

$$m = 1,6749 \times 10^{-24} \text{ g}$$

$$q = 0 \text{ C}$$


Πυρήνας

Ηλεκτρόνιο


αρνητικά φορτισμένο σωματίδιο

$$m = 9,1093897 \times 10^{-28} \text{ g}$$

$$q = 1,6022 \times 10^{-19} \text{ C}$$



Ατομικό μοντέλο Rutherford



Ατομική ακτίνα $\sim 100 \text{ pm} = 1 \times 10^{-10} \text{ m}$

Πυρηνική ακτίνα $\sim 5 \times 10^{-3} \text{ pm} = 5 \times 10^{-15} \text{ m}$

Αν το άτομο είναι στο μέγεθος ενός δωματίου, ο πυρήνας είναι της τάξης ενός κεφαλιού καρφίτσας στο κέντρο του

ΔΟΜΗ ΤΟΥ ΑΤΟΜΟΥ

Ατομικός αριθμός (Z)

αριθμός πρωτονίων του πυρήνα


Μαζικός αριθμός (A)

αριθμός πρωτονίων και νετρονίων του πυρήνα



Ισότοπα

Άτομα των οποίων οι πυρήνες έχουν τον ίδιο ατομικό, αλλά διαφορετικό μαζικό αριθμό



Chadwick's Experiment (1932)

(1935 Noble Prize in Physics)

H atoms - 1 p; He atoms - 2 p

mass He/mass H should = 2

measured mass He/mass H = 4



neutron (n) is neutral (charge = 0)

n mass $\sim$ p mass = 1.67×10^{-24} g

TABLE 2.1 **Mass and Charge of Subatomic Particles**

Particle	Mass (g)	Charge	
		Coulomb	Charge Unit
Electron*	9.10938×10^{-28}	-1.6022×10^{-19}	-1
Proton	1.67262×10^{-24}	$+1.6022 \times 10^{-19}$	+1
Neutron	1.67493×10^{-24}	0	0

*More refined measurements have given us a more accurate value of an electron's mass than Millikan's.

$$\text{mass p} \approx \text{mass n} \approx 1840 \times \text{mass e}^-$$

Atomic number, Mass number and Isotopes


Atomic number (Z) = number of protons in nucleus

Mass number (A) = number of protons + number of neutrons
= atomic number (Z) + number of neutrons

Isotopes are atoms of the same element (X) with different numbers of neutrons in their nuclei



The Isotopes of Hydrogen



How many protons, neutrons, and electrons are in $^{14}_6\text{C}$?

How many protons, neutrons, and electrons are in $^{11}_6\text{C}$?

The Modern Periodic Table



1 1A																	18 8A					
1 H																	2 He					
3																	5 B	6 C	7 N	8 O	9 F	10 Ne
Alkali Metal		Alkali Earth Metal		3 3B	4 4B	5 5B	6 6B	7 7B	8	9	10	11 1B	12 2B	13 Al	Group		15 P	16 S	17 Cl	Noble Gas		
				21 Sc	22 Ti	23 V	24 Cr	25 Mn	26 Fe	27 Co	28 Ni	29 Cu	30 Zn	31 Ga	33 As	34 Se	35 Br					
Alkali Metal		Alkali Earth Metal		39 Y	40 Zr	41 Nb	42 Mo	43 Tc	44 Ru	45 Rh	46 Pd	47 Ag	48 Cd	49 In	Group		51 Sb	52 Te	53 I	Noble Gas		
				55 Cs	56 Ba	57 La	72 Hf	73 Ta	74 W	75 Re	76 Os	77 Ir	78 Pt	79 Au	80 Hg	81 Tl	82 Pb	83 Bi	84 Po	85 At	86 Rn	
Alkali Metal		Alkali Earth Metal		87 Fr	88 Ra	89 Ac	104 Rf	105 Db	106 Sg	107 Bh	108 Hs	109 Mt	110 Ds	111 Rg	112	113	114	115	116	(117)	118	

	Metals
	Metalloids
	Nonmetals


58 Ce	59 Pr	60 Nd	61 Pm	62 Sm	63 Eu	64 Gd	65 Tb	66 Dy	67 Ho	68 Er	69 Tm	70 Yb	71 Lu
90 Th	91 Pa	92 U	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No	103 Lr

Chemistry In Action






Natural abundance of elements in Earth's crust



Natural abundance of elements in human body



Mópio (molecule) is an aggregate of two or more atoms in a definite arrangement held together by chemical forces



A diatomic molecule contains only two atoms

[illegible]

diatomic elements

A ***polyatomic molecule*** contains more than two atoms




An ***ion*** is an atom, or group of atoms, that has a net positive or negative charge.

cation – ion with a positive charge

If a neutral atom **loses** one or more electrons it becomes a cation.




11 protons
11 electrons




11 protons
10 electrons

anion – ion with a negative charge

If a neutral atom **gains** one or more electrons it becomes an anion.




17 protons
17 electrons



17 protons
18 electrons

A ***monatomic ion*** contains only one atom



A ***polyatomic ion*** contains more than one atom











Common Ions Shown on the Periodic Table

1 1A	2 2A											13 3A	14 4A	15 5A	16 6A	17 7A	18 8A
Li ⁺													C ⁴⁻	N ³⁻	O ²⁻	F ⁻	
Na ⁺	Mg ²⁺	3 3B	4 4B	5 5B	6 6B	7 7B	8 8B	9 8B	10 8B	11 1B	12 2B	Al ³⁺		P ³⁻	S ²⁻	Cl ⁻	
K ⁺	Ca ²⁺				Cr ²⁺ Cr ³⁺	Mn ²⁺ Mn ³⁺	Fe ²⁺ Fe ³⁺	Co ²⁺ Co ³⁺	Ni ²⁺ Ni ³⁺	Cu ⁺ Cu ²⁺	Zn ²⁺				Se ²⁻	Br ⁻	
Rb ⁺	Sr ²⁺									Ag ⁺	Cd ²⁺		Sn ²⁺ Sn ⁴⁺		Te ²⁻	I ⁻	
Cs ⁺	Ba ²⁺									Au ⁺ Au ³⁺	Hg ₂ ²⁺ Hg ²⁺		Pb ²⁺ Pb ⁴⁺				

How many protons and electrons are in ${}_{13}^{27}\text{Al}^{3+}$?

How many protons and electrons are in ${}_{34}^{78}\text{Se}^{2-}$?




Formulas and Models

	Hydrogen	Water	Ammonia	Methane
Molecular formula	H_2	H_2O	NH_3	CH_4
Structural formula	$\text{H}-\text{H}$	$\text{H}-\text{O}-\text{H}$	$\begin{array}{c} \text{H}-\text{N}-\text{H} \\ \\ \text{H} \end{array}$	$\begin{array}{c} \text{H} \\ \\ \text{H}-\text{C}-\text{H} \\ \\ \text{H} \end{array}$
Ball-and-stick model				
Space-filling model				



A ***molecular formula*** shows the exact number of atoms of each element in the smallest unit of a substance

An ***empirical formula*** shows the simplest whole-number ratio of the atoms in a substance

molecular






empirical



ionic compounds consist of a combination of cations and an anions

- The formula is usually the same as the empirical formula
- The sum of the charges on the cation(s) and anion(s) in each formula unit must equal zero




The ionic compound NaCl



1A											3A	4A	5A	6A	7A	8A	
	2A																
Li														N	O	F	
Na	Mg											Al			S	Cl	
K	Ca													Br			
Rb	Sr													I			
Cs	Ba																

The most reactive **metals** (green) and the most reactive **nonmetals** (blue) combine to form ionic compounds.

Formula of Ionic Compounds




Chemical Nomenclature

- **Ionic Compounds**

- Often a metal + nonmetal
- Anion (nonmetal), add “ide” to element name



barium chloride



potassium oxide



magnesium hydroxide



potassium nitrate

- Transition metal ionic compounds
 - indicate charge on metal with Roman numerals

A simplified periodic table diagram with the transition metal blocks highlighted in green. The blocks are labeled as follows: 3B, 4B, 5B, 6B, 7B, 8B, 1B, and 2B. The green blocks consist of the 3d and 4d series (3B-8B) and the 5d series (1B-2B).

FeCl_2	2 Cl^- -2 so Fe is +2	iron(II) chloride
FeCl_3	3 Cl^- -3 so Fe is +3	iron(III) chloride
Cr_2S_3	3 S^{2-} -6 so Cr is +3 (6/2)	chromium(III) sulfide

TABLE 2.2**The “-ide” Nomenclature of Some Common Monatomic Anions According to Their Positions in the Periodic Table****Group 4A****Group 5A****Group 6A****Group 7A**C carbide (C^{4-})*N nitride (N^{3-})O oxide (O^{2-})F fluoride (F^-)Si silicide (Si^{4-})P phosphide (P^{3-})S sulfide (S^{2-})Cl chloride (Cl^-)Se selenide (Se^{2-})Br bromide (Br^-)Te telluride (Te^{2-})I iodide (I^-)

*The word “carbide” is also used for the anion C_2^{2-} .

TABLE 2.3

Names and Formulas of Some Common Inorganic Cations and Anions

Cation	Anion
aluminum (Al^{3+})	bromide (Br^-)
ammonium (NH_4^+)	carbonate (CO_3^{2-})
barium (Ba^{2+})	chlorate (ClO_3^-)
cadmium (Cd^{2+})	chloride (Cl^-)
calcium (Ca^{2+})	chromate (CrO_4^{2-})
cesium (Cs^+)	cyanide (CN^-)
chromium(III) or chromic (Cr^{3+})	dichromate ($\text{Cr}_2\text{O}_7^{2-}$)
cobalt(II) or cobaltous (Co^{2+})	dihydrogen phosphate (H_2PO_4^-)
copper(I) or cuprous (Cu^+)	fluoride (F^-)
copper(II) or cupric (Cu^{2+})	hydride (H^-)
hydrogen (H^+)	hydrogen carbonate or bicarbonate (HCO_3^-)
iron(II) or ferrous (Fe^{2+})	hydrogen phosphate (HPO_4^{2-})
iron(III) or ferric (Fe^{3+})	hydrogen sulfate or bisulfate (HSO_4^-)
lead(II) or plumbous (Pb^{2+})	hydroxide (OH^-)
lithium (Li^+)	iodide (I^-)
magnesium (Mg^{2+})	nitrate (NO_3^-)
manganese(II) or manganous (Mn^{2+})	nitride (N^{3-})
mercury(I) or mercurous (Hg_2^{2+})*	nitrite (NO_2^-)
mercury(II) or mercuric (Hg^{2+})	oxide (O^{2-})
potassium (K^+)	permanganate (MnO_4^-)
rubidium (Rb^+)	peroxide (O_2^{2-})
silver (Ag^+)	phosphate (PO_4^{3-})
sodium (Na^+)	sulfate (SO_4^{2-})
strontium (Sr^{2+})	sulfide (S^{2-})
tin(II) or stannous (Sn^{2+})	sulfite (SO_3^{2-})
zinc (Zn^{2+})	thiocyanate (SCN^-)

*Mercury(I) exists as a pair as shown.

- **Molecular compounds**

- Nonmetals or nonmetals + metalloids
- Common names
 - H_2O , NH_3 , CH_4 ,
- Element furthest to the left in a period and closest to the bottom of a group on periodic table is placed first in formula
- If more than one compound can be formed from the same elements, use prefixes to indicate number of each kind of atom
- Last element name ends in *ide*

TABLE 2.4

Greek Prefixes Used in Naming Molecular Compounds

Prefix	Meaning
mono-	1
di-	2
tri-	3
tetra-	4
penta-	5
hexa-	6
hepta-	7
octa-	8
nona-	9
deca-	10

Molecular Compounds

HI hydrogen iodide


NF₃ nitrogen trifluoride

SO₂ sulfur dioxide

N₂Cl₄ dinitrogen tetrachloride

NO₂ nitrogen dioxide


N₂O dinitrogen monoxide



An **acid** can be defined as a substance that yields hydrogen ions (H^+) when dissolved in water.

For example: HCl gas and HCl in water

- Pure substance, hydrogen chloride HCl



- Dissolved in water (H_3O^+ and Cl^-), hydrochloric acid





TABLE 2.5 **Some Simple Acids**

Anion	Corresponding Acid
F^- (fluoride)	HF (hydrofluoric acid)
Cl^- (chloride)	HCl (hydrochloric acid)
Br^- (bromide)	HBr (hydrobromic acid)
I^- (iodide)	HI (hydroiodic acid)
CN^- (cyanide)	HCN (hydrocyanic acid)
S^{2-} (sulfide)	H_2S (hydrosulfuric acid)


An **oxoacid** is an acid that contains hydrogen, oxygen, and another element.




nitric acid




carbonic acid



phosphoric acid



Naming Oxoacids and Oxoanions



The rules for naming **oxoanions**, *anions of oxoacids*, are as follows:

1. When all the H ions are removed from the “-ic” acid, the anion’s name ends with “-ate.”
2. When all the H ions are removed from the “-ous” acid, the anion’s name ends with “-ite.”
3. The names of anions in which one or more but not all the hydrogen ions have been removed must indicate the number of H ions present.

For example:

- H_2PO_4^- dihydrogen phosphate
- HPO_4^{2-} hydrogen phosphate
- PO_4^{3-} phosphate

TABLE 2.6 Names of Oxoacids and Oxoanions That Contain Chlorine

Acid	Anion
HClO_4 (perchloric acid)	ClO_4^- (perchlorate)
HClO_3 (chloric acid)	ClO_3^- (chlorate)
HClO_2 (chlorous acid)	ClO_2^- (chlorite)
HClO (hypochlorous acid)	ClO^- (hypochlorite)

A **base** can be defined as a substance that yields hydroxide ions (OH^-) when dissolved in water.

NaOH	sodium hydroxide
---------------	------------------

KOH	potassium hydroxide
--------------	---------------------

$\text{Ba}(\text{OH})_2$	barium hydroxide
--------------------------	------------------

Hydrates are compounds that have a specific number of water molecules attached to them.

$\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$ barium chloride dihydrate

$\text{LiCl} \cdot \text{H}_2\text{O}$ lithium chloride monohydrate

$\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ magnesium sulfate heptahydrate

$\text{Sr}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$ strontium nitrate tetrahydrate





TABLE 2.7 Common and Systematic Names of Some Compounds


Formula	Common Name	Systematic Name
H ₂ O	Water	Dihydrogen monoxide
NH ₃	Ammonia	Trihydrogen nitride
CO ₂	Dry ice	Solid carbon dioxide
NaCl	Table salt	Sodium chloride
N ₂ O	Laughing gas	Dinitrogen monoxide
CaCO ₃	Marble, chalk, limestone	Calcium carbonate
CaO	Quicklime	Calcium oxide
Ca(OH) ₂	Slaked lime	Calcium hydroxide
NaHCO ₃	Baking soda	Sodium hydrogen carbonate
Na ₂ CO ₃ · 10H ₂ O	Washing soda	Sodium carbonate decahydrate
MgSO ₄ · 7H ₂ O	Epsom salt	Magnesium sulfate heptahydrate
Mg(OH) ₂	Milk of magnesia	Magnesium hydroxide
CaSO ₄ · 2H ₂ O	Gypsum	Calcium sulfate dihydrate

Organic chemistry is the branch of chemistry that deals with carbon compounds


Functional Groups




methanol




methylamine




acetic acid



CH_3OH











CH_3NH_2



CH_3COOH

TABLE 2.8 The First Ten Straight-Chain Alkanes

Name	Formula	Molecular Model
Methane	CH ₄	
Ethane	C ₂ H ₆	
Propane	C ₃ H ₈	
Butane	C ₄ H ₁₀	
Pentane	C ₅ H ₁₂	
Hexane	C ₆ H ₁₄	
Heptane	C ₇ H ₁₆	
Octane	C ₈ H ₁₈	
Nonane	C ₉ H ₂₀	
Decane	C ₁₀ H ₂₂	